



## CHICK STARTER/GROWER PELLET NAB NAP

Complete Chicken Starter/Grower/Finisher/Broiler Feed

### GUARANTEED ANALYSIS

Crude Protein, min	18.0	%	Calcium, max	1.3	%
Lysine, min	0.9	%	Phosphorus, min	0.6	%
Methionine, min	0.3	%	Salt, min	0.2	%
Crude Fat, min	3.0	%	Salt, max	0.7	%
Crude Fiber, max	6.0	%	Phytase (Aspergillus Niger), min	20	SPU/LB*
Calcium, min	0.8	%			

\*One Solid State Fermentation Phytase unit (SPU) is defined as the amount of enzyme that will liberate 1 mmol of inorganic phosphate per minute under the conditions of the assay.

### INGREDIENTS

grain products,processed grain by-products,plant protein products,calcium carbonate,calcium phosphate,salt,choline chloride,vegetable oil,sodium metabisulfite (a preservative),roughage products,hydrolyzed yeast,manganese proteinate,zinc proteinate,hydrated sodium calcium aluminosilicate,dried Aspergillus niger fermentation product,iron proteinate,vitamin E supplement,selenium yeast,mineral oil,niacin supplement,copper proteinate,sodium sulfate,calcium pantothenate,menadione sodium bisulfite complex (source of vitamin K activity),riboflavin supplement,vitamin D-3 supplement,pyridoxine hydrochloride,thiamine mononitrate,natural flavor,vitamin A supplement,ethylenediamine dihydroiodide,vitamin B-12 supplement,folic acid,saccharin sodium,artificial flavor,silicon dioxide,dried Bacillus licheniformis fermentation product,dried Bacillus subtilis fermentation product,biotin,DL-methionine, L-lysine, dried L-lysine fermentation product

**WARNING:** This product, which contains added copper, should not be fed to sheep or related species that have a low tolerance to copper.

This product has been formulated specifically for chickens and is not intended for other species.

**Caution:** Follow label directions. Feeding this feed containing added selenium at levels higher than recommended is not permitted.

### FEEDING DIRECTIONS:

Feed Homestead® Chick Starter/Grower NAB NAP continuously as the sole ration from hatch to desired broiler weight or until 18 weeks of age for laying weight. At 18 weeks of age, transition layer pullets to Homestead Layer®. For poultry flocks with mixed breeds, feed Homestead® All Flock from hatch to maturity or laying age. Provide a constant supply of clean, fresh water.

Hubbard does not use "Restricted-use Proteins" in their products and is in compliance with FDA and state requirements regarding the use, handling and storage of "Restricted-use Protein" products.

### HUBBARD FEEDS

111 West Cherry Street Suite 500  
Mankato, MN 56001  
Net Weight 50 lb (22.68 kg) or bulk invoice.



# 34732

REV 07/22

