



CHICK-EN-EGG® CONCENTRATE PELLET

NAB NAP

Starting/Growing/Laying Chicken Concentrate

GUARANTEED ANALYSIS

Crude Protein, min	38.0	%	Calcium, min	3.5	%
Lysine, min	2.2	%	Calcium, max	4.5	%
Methionine, min	0.8	%	Phosphorus, min	1.6	%
Crude Fat, min	1.0	%	Salt, min	1.0	%
Crude Fiber, max	6.0	%	Salt, max	1.5	%
			Phytase (Aspergillus niger), min	65.0	SPU/LB*

*One Solid State Fermentation Phytase unit (SPU) is defined as the amount of enzyme that will liberate 1 mmol of inorganic phosphate per minute under the conditions of the assay.

INGREDIENTS

plant protein products, calcium carbonate, processed grain by-products, calcium phosphate, salt, vegetable oil, choline chloride, DL-methionine, hydrolyzed yeast, sodium metabisulfite (a preservative), roughage products, manganese proteinate, zinc proteinate, hydrated sodium calcium aluminosilicate, dried Aspergillus niger fermentation product, iron proteinate, vitamin E supplement, ferrous sulfate, selenium yeast, niacin supplement, mineral oil, copper proteinate, calcium pantothenate, menadione sodium bisulfite complex (source of vitamin K activity), riboflavin supplement, sodium sulfate, vitamin D-3 supplement, pyridoxine hydrochloride, thiamine mononitrate, vitamin A supplement, ethylenediamine dihydroiodide, natural flavor, vitamin B-12 supplement, folic acid, saccharin sodium, dried Bacillus licheniformis fermentation product, dried Bacillus subtilis fermentation product, artificial flavor, biotin, silicon dioxide.

WARNING: This product, which contains added copper, should not be fed to sheep or related species that have a low tolerance to copper. This product has been formulated specifically for chickens and is not intended for other species.

Caution: Follow label directions. The addition to feed of higher levels of this concentrate containing added selenium is not permitted.

FEEDING DIRECTIONS:

	Grower	Developer	Layer	Layer
Protein	17%	14%	17%	15%
Corn	1,400	1,550	1,275	1,400
Homestead® Chick-En-Egg® Concentrate NAB NAP	600	450	625	500
Limestone			100	100
TOTAL	2000	2000	2000	2000

Poultry should have access to clean, fresh water at all times. When starting day old birds or after moving or transporting birds, give access to water before putting feed in the feeders. Grit - When birds have access to coarse litter or whole grains, an insoluble grit should be fed. Limit intake of grit to 1 pound per 100 pounds of feed or 2 pounds per 100 birds per week. Grit can be blended with their regular ration or offered free choice in a separate feeder. Do not allow feeders to run empty or stale feed to accumulate. Never feed any feedstuffs that are moldy, musty or suspect in any way.

Hubbard does not use "Restricted-use Proteins" in their products and is in compliance with FDA and state requirements regarding the use, handling and storage of "Restricted-use Protein" products.

HUBBARD FEEDS

111 West Cherry Street Suite 500

Mankato, MN 56001

Net Weight 50 lb (22.68 kg) or bulk invoice.



34740

REV 07/22

